arXiv:math/0308156v2 [math.RT] 18 Sep 2003

Littelmann Paths for the Basic Representation of an Affine Lie Algebra

Peter Magyar

August 13, 2003

Abstract

Let $\mathfrak{g}$ be a complex simple Lie algebra, and $\widehat{\mathfrak{g}}$ the corresponding untwisted affine Lie algebra. Let $\hat{V}(\Lambda_0)$ be the basic irreducible level-one representation of $\widehat{\mathfrak{g}}$, and $\hat{V}_{\lambda^\vee}(\Lambda_0)$ the Demazure module corresponding to the translation $-\lambda^\vee$ in the affine Weyl group. Suppose $\lambda^\vee$ is a sum of minuscule coweights of $\mathfrak{g}$ (which exist if $\mathfrak{g}$ is of classical type or E_6, E_7).

We give a new model for the crystal graphs of $\hat{V}(\Lambda_0)$ and $\hat{V}_{\lambda^\vee}(\Lambda_0)$ which combines Littelmann's path model and the Kyoto path model. As a corollary, we prove that $V_{\lambda^\vee}(\Lambda_0)$ is isomorphic as a $\mathfrak{g}$-module to a tensor product of fundamental representations of $\mathfrak{g}$.

1 Main Results

1.1 Product Theorems

Let $\mathfrak{g}$ be a complex simple Lie algebra and $\widehat{\mathfrak{g}}$ the corresponding untwisted affine Kac-Moody algebra. The basic representation $\hat{V}(\Lambda_0)$ is the simplest and most important $\widehat{\mathfrak{g}}$-module (see Sec. 2.1 for definitions, as well as [6, Ch. 14],[21, Ch. 10]). One of its remarkable properties is the Tensor Product Phenomenon. In many cases, the Demazure modules $\hat{V}_z(\Lambda_0) \subset \hat{V}(\Lambda_0)$ are representations of the finite-dimensional algebra $\mathfrak{g}$, and they factor into a tensor product of many small $\mathfrak{g}$-modules. Hence the full $\hat{V}(\Lambda_0)$ could be constructed by extending the $\mathfrak{g}$-structure on the semi-infinite tensor power $V \otimes V \otimes \cdots$ of a small $\mathfrak{g}$-module V.

The Kyoto school of Jimbo, Kashiwara et al., has established this phenomenon for $\mathfrak{g}$ of classical type (and for a large class of $\widehat{\mathfrak{g}}$-modules $\hat{V}(\Lambda)$) via the theory of perfect crystals [7],[10],[11],[4],[9] a development of the earlier theory of semi-infinite paths [2]. Pappas and Rapoport [20] have given a geometric version of the phenomenon for type A: they construct a flat deformation of Schubert varieties of the affine Grassmannian into a product of finite Grassmannians.

In this paper, we extend the Tensor Product Phenomenon for $\hat{V}(\Lambda_0)$ to the non-classical types E_6 and E_7 by a uniform method which applies whenever $\mathfrak{g}$ possesses a minuscule representation, or more precisely a minuscule coweight. We shall rely on a key property of such coweights which may be taken as the definition. Let $\hat{X}$ be the extended Dynkin diagram (the diagram of $\widehat{\mathfrak{g}}$). A coweight $\varpi^\vee$ of $\mathfrak{g}$ is *minuscule* if and only if it is a fundamental coweight $\varpi^\vee = \varpi_i^\vee$ and there exists an automorphism σ of $\hat{X}$ taking the node i to the distinguished node 0. Such automorphisms exist in types A, B, C, D, E_6, E_7.

We let $V(\lambda)$ denote the irreducible $\mathfrak{g}$-module with highest weight λ, and $V(\lambda)^*$ its dual module. Our main representation-theoretic result is:

Theorem 1 *Let $\lambda^\vee$ be an element of the coroot lattice of $\mathfrak{g}$ which is a sum:*

$$\lambda^\vee = \lambda_1^\vee + \cdots + \lambda_m^\vee,$$

where $\lambda_1^\vee, \cdots, \lambda_m^\vee$ are minuscule fundamental coweights (not necessarily distinct), with corresponding fundamental weights $\lambda_1, \cdots, \lambda_m$. Let $\hat{V}_{\lambda^\vee}(\Lambda_0) \subset \hat{V}(\Lambda_0)$ be the Demazure module corresponding to the anti-dominant translation $t_{-\lambda^\vee}$ in the affine Weyl group.

Then there is an isomorphism of $\mathfrak{g}$-modules:

$$\hat{V}_{\lambda^\vee}(\Lambda_0) \cong V(\lambda_1)^* \otimes \cdots \otimes V(\lambda_m)^* .$$

Now fix a minuscule coweight $\varpi^\vee$ and its corresponding fundamental weight ϖ. Let N be the smallest positive integer such that $N\varpi^\vee$ lies in the coroot lattice of $\mathfrak{g}$. Then we have the following characterization of the basic irreducible $\widehat{\mathfrak{g}}$-module:

Theorem 2 *The tensor power $V_N := V(\varpi)^{\otimes N}$ possesses non-zero $\mathfrak{g}$-invariant vectors. Fix such a vector v_N, and define the $\mathfrak{g}$-module $V^{\otimes\infty}$ as the direct limit of the sequence:*

$$V_N \hookrightarrow V_N^{\otimes 2} \hookrightarrow V_N^{\otimes 3} \hookrightarrow \cdots$$

where each inclusion is defined by: $v \mapsto v_N \otimes v$.

Then $\hat{V}(\Lambda_0)$ is isomorphic as a $\mathfrak{g}$-module to $V^{\otimes\infty}$.

It would be interesting to define the action of the full algebra $\widehat{\mathfrak{g}}$ on $V^{\otimes\infty}$, and thus give a uniform "path construction" of the basic representation (cf. [2]): that is, to define the raising and lowering operators E_0, F_0, as well as the charge operator d. Combinatorial definitions of the charge produce for $\mathfrak{g}$ of classical type produce generalizations of the Kostka-Foulkes polynomials (c.f. [19]).

1.2 Crystal Theorems

Our basic tool to prove the above results is Littelmann's combinatorial model [15],[16],[14] for representations of Kac-Moody algebras, a vast generalization of Young tableaux. Littelmann's paths and path operators give a flexible construction of the crystal graphs associated to quantum $\mathfrak{g}$-modules by Kashiwara [8] and Lusztig [17] (see also [5],[4]). Roughly speaking, we prove Theorem 1 by reducing it to an identity of paths: we construct a path crystal for the affine Demazure module which is at the same time a path crystal for the tensor product. We carry this out in Sec. 2.

Theorem 2 follows as a corollary in Sec. 3. To describe the crystal graph of the semi-infinite tensor product, we pass to a semi-infinite limit of Littelmann paths which we call *skeins*. We thus recover the Kyoto path model for classical $\mathfrak{g}$, and our results are equally valid for E_6, E_7.

To be more precise, we briefly sketch Littelmann's theory. Let us define a $\mathfrak{g}$-*crystal* as a *set* $\mathcal{B}$ with a *weight function* $\mathrm{wt} : \mathcal{B} \to \oplus_{i=1}^{r} \mathbb{Z}\varpi_i$, and partially defined *crystal operators* $e_1, \dots, e_r, f_1, \dots, f_r : \mathcal{B} \to \mathcal{B}$ satisfying:

$$\mathrm{wt}(f_i(b)) = \mathrm{wt}(b) - \alpha_i \qquad \text{and} \qquad e_i(b){=}b' \Longleftrightarrow f_i(b'){=}b\,.$$

Here $\varpi_1, \dots, \varpi_r$ are the fundamental weights and $\alpha_1, \dots, \alpha_r$ are the roots of $\mathfrak{g}$. A *dominant* element is a $b{\in}B$ such that $e_i(b)$ is not defined for any i. We say that a crystal $\mathcal{B}$ is a *model* for a $\mathfrak{g}$-module V if the formal character of $\mathcal{B}$ is equal to the character of V, and the dominant elements of $\mathcal{B}$ correspond to the highest-weight vectors of V. That is:

$$\mathrm{char}(V) = \textstyle\sum_{b \in B} e^{\mathrm{wt}(b)} \qquad \text{and} \qquad V \cong \bigoplus_{b \text{ dom}} V(\mathrm{wt}(b))\,,$$

where the second sum is over the dominant elements of $\mathcal{B}$. Clearly, a $\mathfrak{g}$-module V is determined up to isomorphism by any model $\mathcal{B}$.

We construct such $\mathfrak{g}$-crystals $\mathcal{B}$ consisting of polygonal paths in the vector space of weights, $\mathfrak{h}^*_{\mathbb{R}} := \oplus_{i=1}^{r} \mathbb{R}\varpi_i$. Specifically:

- The *elements* of $\mathcal{B}$ are continuous piecewise-linear mappings $\pi : [0,1] \to \mathfrak{h}^*_{\mathbb{R}}$, up to reparametrization, with initial point $\pi(0) = 0$. We use the notation $\pi = (v_1 \star v_2 \star \cdots \star v_k)$, where $v_1, \dots, v_k \in \mathfrak{h}^*_{\mathbb{R}}$ are vectors, to denote the polygonal path starting at 0 and moving linearly to v_1, then to $v_1{+}v_2$, etc.
- The *weight* of a path is its endpoint:

  $$\mathrm{wt}(\pi) := \pi(1) = v_1{+}\cdots{+}v_k\,.$$

- The *crystal lowering operator* f_i is defined as follows (and there is a similar definition of the raising operator e_i). Let $\star$ denote the natural associative operation of concatenation of paths, and let any linear map $w : \mathfrak{h}^*_{\mathbb{R}} \to \mathfrak{h}^*_{\mathbb{R}}$ act pointwise on paths: $w(\pi) := (w(v_1) \star \cdots \star w(v_k))$. We will divide a path π into three well-defined sub-paths, $\pi = \pi_1 \star \pi_2 \star \pi_3$, and reflect the middle piece by the simple reflection s_i:

  $$f_i\pi := \pi_1 \star s_i\pi_2 \star \pi_3\,.$$

> The pieces π_1, π_2, π_3 are determined according to the behavior of the i-height function $h_i(t) = h_i^\pi(t) := \langle \pi(t), \alpha_i^\vee \rangle$, as the point $\pi(t)$ moves along the path from $\pi(0) = 0$ to $\pi(1) = \mathrm{wt}(\pi)$. This function may attain its minimum value $h_i(t) = M$ several times. If, after the *last* minimum point, $h_i(t)$ never rises to the value $M{+}1$, then $f_i\pi$ is *undefined.* Otherwise, we define π_2 as the last sub-path of π on which $M \leq h_i(t) \leq M{+}1$, and π_1, π_3 as the remaining initial and final pieces of π.

A key advantage of the path model is that the crystal operators, while complicated, are universally defined for all paths. Hence a path crystal is completely specified by giving its set of paths $\mathcal{B}$.

Also, the dominant elements have a neat pictorial characterization, as the paths π which never leave the fundamental Weyl chamber: that is, $\mathfrak{h}_i^\pi(t) \geq 0$ for all $t \in [0,1]$ and all $i = 1, \ldots, r$. For simplicity we restrict ourselves to *integral* dominant paths, meaning that all the steps are integral weights: $v_1, \ldots, v_k \in \oplus_{i=1}^r \mathbb{Z}\varpi_i$. (For arbitrary dominant paths, see [16].)

Littelmann's Character Theorem [16] states that if π is any integral dominant path with weight λ, then the set of paths $\mathcal{B}(\pi)$ generated from π by $f_1, \ldots, f_r$ is a model for the irreducible $\mathfrak{g}$-module $V(\lambda)$. (This $\mathcal{B}(\pi)$ is also closed under $e_1, \ldots, e_r$.) Note that we can choose *any* integral path π which stays within the Weyl chamber and ends at λ, and each such choice gives a different (but isomorphic) path crystal modelling $V(\lambda)$. In principle, any reasonable indexing set for a basis of $V(\lambda)$ should be in natural bijection with $\mathcal{B}(\pi)$ for some choice of π. For example, classical Young tableaux correspond to choosing the steps v_j to be coordinate vectors in $\mathfrak{h}_\mathbb{R}^* \cong \mathbb{R}^n$.

Furthermore, we have Littelmann's Product Theorem [16]: if $\pi_1, \ldots, \pi_m$ are dominant integral paths of respective weight $\lambda_1, \ldots, \lambda_m$, then $\mathcal{B}(\pi_1) \star \cdots \star \mathcal{B}(\pi_m)$, the set of all concatenations, is a model for the tensor product $V(\lambda_1) \otimes \cdots \otimes V(\lambda_m)$.

Everything we have said also holds for the affine algebra $\widehat{\mathfrak{g}}$, provided we replace the roots $\alpha_1, \ldots, \alpha_r$ of $\mathfrak{g}$ by the roots $\alpha_0, \alpha_1, \ldots, \alpha_r$ of $\widehat{\mathfrak{g}}$; and the weights $\varpi_1, \ldots, \varpi_r$ of $\mathfrak{g}$ by the weights $\Lambda_0, \Lambda_1, \ldots, \Lambda_r$ of $\widehat{\mathfrak{g}}$. We also replace the vector space $\mathfrak{h}_\mathbb{R}^*$ by $\widehat{\mathfrak{h}}_\mathbb{R}^* := \oplus_{i=0}^r \mathbb{R}\Lambda_i \oplus \mathbb{R}\delta$, where δ is the non-divisible positive imaginary root of $\widehat{\mathfrak{g}}$. (Indeed, the theory works uniformly for all symmetrizable Kac-Moody algebras.) We denote path crystals for $\mathfrak{g}$ and $\widehat{\mathfrak{g}}$ by $\mathcal{B}$ and $\hat{\mathcal{B}}$ respectively.

We can also model the affine Demazure module $\hat{V}_z(\Lambda)$, where $z \in \widehat{W}$, the Weyl group of $\widehat{\mathfrak{g}}$. Indeed, if $z = s_{i_1} \cdots s_{i_m}$ is a reduced decomposition and π is an integral dominant path of weight Λ, we define the *Demazure path crystal*:

$$\hat{\mathcal{B}}_z(\pi) := \{ f_{i_1}^{k_1} \cdots f_{i_m}^{k_m} \pi \mid k_1, \ldots, k_m \geq 0 \} \, .$$

Then the formal character of $\hat{\mathcal{B}}_z(\pi)$ is equal to the character of $\hat{V}_z(\Lambda)$, and π is the unique dominant path [15]. Now suppose $z = t_{-\lambda^\vee}$, an anti-dominant translation in $\widehat{W}$, so that $\hat{V}_{\lambda^\vee}(\Lambda) := \hat{V}_z(\Lambda)$ is a $\mathfrak{g}$-submodule of $\hat{V}(\Lambda)$; and consider $\hat{\mathcal{B}}_{\lambda^\vee}(\pi) := \hat{\mathcal{B}}_z(\pi)$ as a $\mathfrak{g}$-crystal by forgetting the action of f_0, e_0 and projecting the affine weight function to $\mathfrak{h}^*_{\mathbb{R}}$. Then Littelmann's Restriction Theorem [16] implies that the $\mathfrak{g}$-crystal $\hat{\mathcal{B}}_{\lambda^\vee}(\pi)$ is a model for the $\mathfrak{g}$-module $\hat{V}_{\lambda^\vee}(\Lambda)$.

Now we are ready to state our main combinatorial results. For λ a dominant weight, define its dual weight λ^* by the dual $\mathfrak{g}$-module: $V(\lambda^*) = V(\lambda)^*$.

Theorem 3 *Let $\lambda^\vee$ be as in Theorem 1. Then there exists an integral $\widehat{\mathfrak{g}}$-dominant path π with weight Λ_0 which generates the $\widehat{\mathfrak{g}}$ Demazure path crystal:*

$$\hat{\mathcal{B}}_{\lambda^\vee}(\pi) = \Lambda_0 \star \mathcal{B}(\lambda_1^*) \star \cdots \star \mathcal{B}(\lambda_m^*) \mod \mathbb{R}\delta\,.$$

This is to be understood as an equality of sets of paths in $\widehat{\mathfrak{h}}^*_{\mathbb{R}}$ mod $\mathbb{R}\delta$.

Theorem 1 follows immediately from this. Indeed, $s_i\Lambda_0 = \Lambda_0$ for $i = 1, \ldots, r$, so $f_i(\Lambda_0 \star \pi') = \Lambda_0 \star f_i(\pi')$ for any path π'. Thus the crystal on the right-hand side of the above equation is isomorphic to $\mathcal{B}(\lambda_1^*) \star \cdots \star \mathcal{B}(\lambda_m^*)$, which models $V(\lambda_1)^* \otimes \cdots \otimes V(\lambda_r)^*$. See [3] for methods of enumerating the paths in this crystal (and hence computing the dimension of the corresponding representation).

Next we give the following crystal version of Theorem 2:

Theorem 4 *Let $\varpi^\vee$, N be as in Theorem 2. Define the N-fold concatenation $\mathcal{B}_N = \mathcal{B}(\varpi^*) \star \cdots \star \mathcal{B}(\varpi^*)$. Then $\Lambda_0 \star \mathcal{B}_N$ contains a unique $\widehat{\mathfrak{g}}$-dominant path $\Lambda_0 \star \pi_N$.*

Define $\hat{\mathcal{B}}_\infty$ as the direct limit of the sequence:

$$\Lambda_0 \star \mathcal{B}_N \hookrightarrow \Lambda_0 \star \mathcal{B}_N \star \mathcal{B}_N \hookrightarrow \Lambda_0 \star \mathcal{B}_N \star \mathcal{B}_N \star \mathcal{B}_N \hookrightarrow \cdots$$

where the inclusions are given by $\Lambda_0 \star \pi \mapsto \Lambda_0 \star \pi_N \star \pi$. Then $\hat{\mathcal{B}}_\infty$ has a natural $\widehat{\mathfrak{g}}$-crystal structure which is isomorphic to the $\widehat{\mathfrak{g}}$-crystal of $\hat{V}(\Lambda_0)$.

In Sec. 3.2, we will use crystals of semi-infinite paths ("skeins") to give a natural meaning to the direct limit above.

1.3 Example: E_6

Referring to Bourbaki [1], we write the extended Dynkin diagram $\hat{X} = \hat{E}_6$:



The simple roots are defined inside $\mathbb{R}^6$ with standard basis $\epsilon_1, \ldots, \epsilon_6$. (Our ϵ_6 is $\frac{1}{\sqrt{3}}(-\epsilon_6 - \epsilon_7 + \epsilon_8)$ in Bourbaki's notation.) They are:

$$\alpha_1 = \tfrac{1}{2}(\epsilon_1 + \epsilon_2 + \epsilon_3 + \epsilon_4 + \epsilon_5) + \tfrac{\sqrt{3}}{2}\epsilon_6, \quad \alpha_2 = \epsilon_1 + \epsilon_2,$$
$$\alpha_3 = \epsilon_2 - \epsilon_1, \quad \alpha_4 = \epsilon_3 - \epsilon_2, \quad \alpha_5 = \epsilon_4 - \epsilon_3, \quad \alpha_6 = \epsilon_5 - \epsilon_4\,.$$

Since E_6 is simply laced, the coroots and coweights may be identified with the roots and weights, with the natural pairing given by the standard dot product on $\mathbb{R}^6$.

We focus on the minuscule coweight $\varpi_1^\vee$ corresponding to the diagram automorphism σ with $\sigma(1) = 0$ and $\sigma(0) = 6$. In this case, the corresponding fundamental representation $V(\varpi_1)$ is also minuscule, meaning that all of its weights are extremal weights $\lambda \in W(E_6){\cdot}\varpi_1$. The roots $\alpha_2, \cdots, \alpha_6$ generate the root sub-system $D_5 \subset E_6$, and the reflection subgroup $W(D_5) = \mathrm{Stab}_{W(E_6)}(\varpi_1)$ acts by permuting $\epsilon_1, \ldots, \epsilon_5$ (the subgroup $W(A_4) = S_5$) and by changing an even number of signs $\pm\epsilon_1, \ldots, \pm\epsilon_5$. We have $\dim V(\varpi_1) = |W(E_6)/W(D_5)| = 27$. The weights are:

$$
\begin{gathered}
\varpi_1 = \tfrac{2\sqrt{3}}{3}\epsilon_6, \\
S_5{\cdot}\tfrac{1}{2}(-\epsilon_1+\epsilon_2+\epsilon_3+\epsilon_4+\epsilon_5) + \tfrac{\sqrt{3}}{6}\epsilon_6, \\
S_5{\cdot}\tfrac{1}{2}(-\epsilon_1-\epsilon_2-\epsilon_3+\epsilon_4+\epsilon_5) + \tfrac{\sqrt{3}}{6}\epsilon_6, \\
-\tfrac{1}{2}(\epsilon_1+\epsilon_2+\epsilon_3+\epsilon_4+\epsilon_5) + \tfrac{\sqrt{3}}{6}\epsilon_6, \\
\pm S_5{\cdot}\epsilon_1 - \tfrac{\sqrt{3}}{3}\epsilon_6,
\end{gathered}
$$

The lowest weight is $-\varpi_6 = -\epsilon_5 - \frac{\sqrt{3}}{3}\epsilon_6$, so that $V(\varpi_1)^* = V(\varpi_6)$ and $\varpi_1^* = \varpi_6$.

The simplest path crystal for $V(\varpi_1^*)$ is the set of 27 straight-line paths from 0 to the negatives of the above extremal weights:

$$\mathcal{B}(\varpi_1^*) = \{\, (v) \mid v \in -W(E_6){\cdot}\varpi_1 \,\}$$

We have $3\varpi_1^\vee \in \oplus_{i=1}^6 \mathbb{R}\alpha_i^\vee$ the coroot lattice, so that $N = 3$ in Theorem 2, and this N is also the order of the automorphism σ. The path crystal $\mathcal{B}_3 := \mathcal{B}(\varpi_1^*) \star \mathcal{B}(\varpi_1^*) \star \mathcal{B}(\varpi_1^*)$, the set of all 3-step walks with steps chosen from the 27 weights of $V(\varpi_1^*)$, is a model for $V(\varpi_1^*)^{\otimes 3}$. In this case there is a unique $\mathfrak{g}$-dominant path of weight 0,

$$\pi_3 := (\varpi_6) \star (\varpi_1 - \varpi_6) \star (-\varpi_1)\,,$$

which corresponds to the one-dimensional space of $\mathfrak{g}$-invariant vectors in $V(\varpi_i^*)^{\otimes 3}$.

Now Theorem 3 states that the affine Demazure module $\hat{V}_{3m\varpi_1^\vee}(\Lambda_0)$ is modelled by the $\widehat{\mathfrak{g}}$-path crystal:

$$\mathcal{B}_{3m} = \{(\Lambda_0 \star v_1 \star \cdots \star v_{3m}) \mid v_j \in -W(E_6){\cdot}\varpi_1\}\,,$$

the set of all $3m$-step walks in $\Lambda_0 \oplus \mathbb{R}^6$ starting at Λ_0, with steps chosen from the 27 weights of $V(\varpi_1^*)$. This path crystal is generated from its unique $\widehat{\mathfrak{g}}$-dominant path $\Lambda_0 \star \pi_3 \star \cdots \star \pi_3$. As a corollary of Theorem 4, we can realize the $\widehat{\mathfrak{g}}$-crystal of the basic $\widehat{\mathfrak{g}}$-module $\hat{V}(\Lambda_0)$ as the set of all infinite walks (or "skeins") of the form:

$$\pi = \Lambda_0 \star \underbrace{\pi_3 \star \cdots \star \pi_3}_{\text{infinite}} \star v_1 \star \cdots \star v_{3m}\,,$$

with $m > 0$ and $v_j \in -W(E_6)\cdot\varpi_1$. The endpoint of such a skein is $\mathrm{wt}(\pi) := \Lambda_0+v_1+\cdots+v_m$. The crystal operators f_i are defined just as for finite paths. Acting near the end of the skein, they unwind the coils π_3 one at a time, right-to-left. See Sec. 3 for details.

2 Demazure Crystals

2.1 Notations

We will work with a complex simple Lie algebra $\mathfrak{g}$ of rank r, a Cartan subalgebra $\mathfrak{h} \subset \mathfrak{g}$, the set of roots $\Delta \subset \mathfrak{h}^*$, and the set of coroots $\Delta^\vee \subset \mathfrak{h}$. We write the highest root of Δ as $\theta = a_1\alpha_1 + \cdots + a_r\alpha_r$, and its coroot as $\theta^\vee = a_1^\vee\alpha_1^\vee + \cdots + a_r^\vee\alpha_r^\vee$. Warning: If $\mathfrak{g}$ is not simply laced, $\theta^\vee$ is not the highest root of the dual root system $\Delta^\vee$.

The Weyl group W of $\mathfrak{g}$ is generated by reflections $s_1, \ldots, s_r$ defined by $s_i(\lambda) = \lambda - \langle\lambda, \alpha_i^\vee\rangle\alpha_i$ for $\lambda \in \mathfrak{h}^*_{\mathbb{R}} := \oplus_{i=1}^r \mathbb{R}\varpi_i$. We have the fundamental Weyl chamber $C = \{\lambda \in \mathfrak{h}^*_{\mathbb{R}} \mid \langle\lambda, \alpha_i^\vee\rangle \geq 0, \quad i{=}1, \ldots, r\}$. The Weyl group also acts naturally on $\mathfrak{h}_{\mathbb{R}}$. If we choose a W-invariant bilinear form $(\,\cdot\,|\,\cdot\,)$ on $\mathfrak{h}_{\mathbb{R}}$, we have the isomorphism $\nu : \mathfrak{h}_{\mathbb{R}} \to \mathfrak{h}^*_{\mathbb{R}}$ defined by $\langle\nu(h), h'\rangle = (h|h')$ for $h, h' \in \mathfrak{h}_{\mathbb{R}}$. We normalize so that $\nu(\theta^\vee) = \theta$ and $\nu(\varpi_i^\vee) = \frac{a_i}{a_i^\vee}\varpi_i$.

Now let $\widehat{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}K \oplus \mathbb{C}d$ be the untwisted affine Lie algebra of $\mathfrak{g}$, where K is a central element and $d = t\frac{d}{dt}$ is a derivation. (Cf. Kac [6, Ch. 6 and 7].) Then $\widehat{\mathfrak{g}}$ has Cartan subalgebra $\widehat{\mathfrak{h}} = \mathfrak{h} \oplus \mathbb{C}K \oplus \mathbb{C}d$, with dual $\widehat{\mathfrak{h}}^* = \mathfrak{h}^* \oplus \mathbb{C}\Lambda_0 \oplus \mathbb{C}\delta$, where $\langle\Lambda_0, \mathfrak{h}\rangle = \langle\delta, \mathfrak{h}\rangle = 0$ and $\langle\Lambda_0, K\rangle = \langle\delta, d\rangle = 1$.

The simple roots of $\widehat{\mathfrak{g}}$ are $\alpha_1, \ldots, \alpha_r$ and $\alpha_0 = \delta{-}\theta$; the simple coroots are $\alpha_1^\vee, \ldots, \alpha_r^\vee$ and $\alpha_0^\vee = K{-}\theta^\vee$. The fundamental weights are Λ_0 and $\Lambda_i = \varpi_i + a_i^\vee\Lambda_0$ for $i = 1, \ldots, r$. The affine Weyl group $\widehat{W}$ is generated by the reflections $s_0, s_1, \ldots, s_r$ acting on $\widehat{\mathfrak{h}}^*_{\mathbb{R}}$. The fundamental Weyl chamber of $\widehat{\mathfrak{g}}$ is the cone $\hat{C} = \{\Lambda \in \widehat{\mathfrak{h}}^*_{\mathbb{R}} \mid \langle\Lambda, \alpha_i^\vee\rangle \geq 0,\ i{=}0, \ldots, r\}$ with extremal rays $\Lambda_0, \ldots, \Lambda_r$.

For $\Lambda \in \oplus_{i=0}^r \mathbb{N}\Lambda_i$ and $w \in \widehat{W}$, we have the irreducible highest-weight $\widehat{\mathfrak{g}}$-module $\hat{V}(\Lambda)$. We will also consider the Demazure module $\hat{V}_z(\Lambda) := U(\widehat{\mathfrak{n}}_+)\cdot v_{z\Lambda}$, where $\widehat{\mathfrak{n}}_+$ is the algebra spanned by the positive weight-spaces of $\widehat{\mathfrak{g}}$, $z \in \widehat{W}$ is a Weyl group element, and $v_{z\Lambda}$ is a non-zero vector of extremal weight $z\Lambda$ in $\hat{V}(\Lambda)$.

For a weight $\Lambda = \lambda + \ell\Lambda_0 + m\delta$, we will ignore the charge $m = \langle\Lambda, d\rangle$ and work modulo $\mathbb{R}\delta$, even when we do not indicate this explicitly. Since $\langle\delta, \alpha_i^\vee\rangle = 0$ for $i = 0, \ldots, r$, the charge has no effect on the path operators f_i, e_i.

Consider the lattice $M = \nu(\oplus_{i=1}^r \mathbb{Z}\alpha_i^\vee)$ in $\mathfrak{h}^*_{\mathbb{R}}$. For any $\mu \in M$, there is an element $t_\mu \in \widehat{W}$ which acts on the weights of level ℓ as translation by $\ell\mu$: that is,

$$t_\mu(\lambda{+}\ell\Lambda_0) = \lambda{+}\ell\mu{+}\ell\Lambda_0 \pmod{\mathbb{R}\delta}.$$

Furthermore, the affine Weyl group is a semi-direct product of the finite Weyl group with the lattice of translations: $\widehat{W} = W \ltimes t_M$.

Consider the anti-dominant translation $z = t_{-\lambda}$ corresponding to a dominant weight $\lambda = \nu(\lambda^\vee) \in C \cap M$. We denote the resulting Demazure module as $\hat{V}_{\lambda^\vee}(\Lambda_0) := \hat{V}_z(\Lambda_0)$. Then the $\widehat{\mathfrak{n}}_+$-module $\hat{V}_{\lambda^\vee}(\Lambda_0)$ is also a $\mathfrak{g}$-submodule of $\hat{V}(\Lambda_0)$:

$$\mathfrak{g}\cdot\hat{V}_{\lambda^\vee}(\Lambda_0) \subset \hat{V}_{\lambda^\vee}(\Lambda_0)\,,$$

and these are the only $z \in \widehat{W}$ for which $\hat{V}_z(\Lambda_0)$ is a $\mathfrak{g}$-module.

2.2 Minuscule weights and coweights

We collect needed facts concerning minuscule weights in root systems. The statements below are well-known and easily verified from tables [1],[6], although direct proofs are also not difficult (cf. [18]).

We say a non-zero coweight $\varpi^\vee \in \mathfrak{h}^*_{\mathbb{R}}$ is *minuscule for* Δ if $\langle \alpha, \varpi^\vee \rangle = 0$ or 1 for all positive roots $\alpha \in \Delta_+$. Equivalently, $\varpi^\vee = \varpi_i^\vee$ for some $i = 1, \ldots, r$ with $a_i = \langle \theta, \varpi_i^\vee \rangle = 1$. This implies that $a_i^\vee = 1$ as well, so that $\nu(\varpi_i^\vee) = \varpi_i$. The classification of the minuscule $\varpi_i^\vee$ is most concisely described by listing the pairs $(X,\, X\backslash\{i\})$, where X is the Dynkin diagram of $\mathfrak{g}$. We have $\varpi_i^\vee$ minuscule when:

$$\begin{array}{rcl}(X,\, X\backslash\{i\}) & \cong & (A_r, A_{r-k}{\times}A_{k-1}),\quad k{=}1,\ldots,r \\ & & (B_r, B_{r-1}),\quad (C_r, A_{r-1}), \\ & & (D_r, D_{r-1}),\quad (D_r, A_r), \\ & & (E_6, D_5),\quad (E_7, E_6)\,. \end{array}$$

There are no minuscule $\varpi_i^\vee$ for $X = E_8, F_4$, or G_2.

Now define the extended Weyl group $\widetilde{W}$ as a group of linear mappings on $\widehat{\mathfrak{h}}^*_{\mathbb{R}}$: namely, $\widetilde{W} := W \ltimes t_L$, where $L = \nu(\oplus_{i=1}^r \mathbb{Z}\varpi_i^\vee)$. Let

$$\Sigma := \{\sigma \in \widetilde{W} \mid \sigma(\hat{C}) = \hat{C}\}\,,$$

the symmetries in $\widetilde{W}$ of the fundamental chamber of $\widehat{\mathfrak{h}}^*_{\mathbb{R}}$. The set Σ is a system of coset representatives for $\widetilde{W}/\widehat{W}$, so that $\widetilde{W} = \Sigma \ltimes \widehat{W}$. We can extend the Bruhat length function to $\widetilde{W}$ as: $l(\sigma w) = l(w\sigma) := l(w)$ for $\sigma \in \Sigma$, $w \in \widehat{W}$. Each element $\sigma \in \Sigma$ defines an automorphism of the Dynkin diagram of $\widehat{\mathfrak{g}}$ which we also write as σ. For $j = 0, \ldots, r$, we have:

$$\sigma(\Lambda_j) = \Lambda_{\sigma(j)} \quad \text{and} \quad \sigma(\alpha_j) = \alpha_{\sigma(j)}\,.$$

There is a natural correspondence between elements of Σ and minuscule coweights. Each $\sigma \in \Sigma$ can be written uniquely as:

$$\sigma = \bar{\sigma}\, t_{-\nu(\varpi_i^\vee)} = \bar{\sigma}\, t_{-\varpi_i}\,,$$

for $\bar{\sigma} \in W$ and $\varpi_i^\vee$ a minuscule coweight. In fact, $\bar{\sigma} = w_0 w_i$, where w_0 is the longest element of W and w_i is the longest element of the parabolic subgroup $W_i := \mathrm{Stab}_W(\varpi_i)$. We have $\bar{\sigma}(\alpha_j) = \alpha_{\sigma(j)}$ for $j \neq i$, and $\bar{\sigma}(\alpha_i) = -\theta$.

We have $\sigma(\Lambda_i) = \bar{\sigma} t_{-\varpi_i}(\varpi_i + \Lambda_0) = \Lambda_0$, so that

$$\sigma(i) = 0\,.$$

Also, $\bar{\sigma}(\varpi_i) = w_0 w_i(\varpi_i) = w_0(\varpi_i) = -\varpi_i^*$, and $\varpi_{\sigma(0)} = \varpi_i^*$.

The number N appearing in Theorems 2 and 4 is the order of σ in the group Σ, and is also the order of $\varpi^\vee$ in $\oplus_{i=1}^r \mathbb{R}\varpi_i^\vee \,/\, \oplus_{i=1}^r \mathbb{R}\alpha_i^\vee$.

The definition of a minuscule weight ϖ for $\Delta^\vee$ is dual to the above: $\langle \varpi, \alpha^\vee \rangle = 0$ or 1 for all positive coroots $\alpha^\vee \in \Delta_+^\vee$; or equivalently $\varpi = \varpi_i$ and $\langle \varpi, \theta^* \rangle = 1$, where θ^* is highest in the root system $\Delta^\vee$. The fundamental representation $V(\varpi)$ corresponding to a minuscule ϖ has a basis consisting of extremal weight vectors $v_{w(\varpi)}$ for $w \in W$. Note, however, that ϖ need not be minuscule even when the corresponding coweight $\varpi^\vee$ is minuscule. In fact, we have:

Lemma 5 *Let $\varpi_i^\vee$ be a minuscule coweight for Δ, and ϖ_i the corresponding weight.*

(i) If Δ is simply laced (i.e., all root vectors have the same length), then ϖ_i is minuscule for $\Delta^\vee$.

(ii) If Δ is not simply laced, then ϖ_i is minuscule for $\Delta_s^\vee$, the simply-laced root system of short vectors in $\Delta^\vee$. Furthermore, $\alpha_i^\vee \in \Delta_s^\vee$.

Proof. Part (i) follows from $\nu(\varpi_i^\vee) = (a_i/a_i^\vee)\,\varpi_i = \varpi_i$. Part (ii) is immediately verified for the relevant types B_r and C_r. □

Consider the *parabolic Bruhat order* on the W-orbit $W\cdot\varpi_i$. That is, the partial order generated by the relations: $\tau_1 < \tau_2$ if $\tau_1 = \tau_2 - d\alpha$ for some positive root α and some $d > 0$. The following result says that if $\varpi_i^\vee$ is minuscule, then this "strong" order is identical to the "weak" order:

Lemma 6 (Stembridge [22]) *Suppose the coweight $\varpi_i^\vee$ is minuscule. Then the Bruhat order on $W\cdot\varpi_i$ has covering relations:*

$$\tau_1 \lessdot \tau_2 \quad \textit{whenever} \quad \tau_1 = \tau_2 - d\alpha_j$$

for some simple *root α_j of W and some $d > 0$.*

Proof. This follows from Stembridge's formulation by noting that $W\cdot\varpi_i \cong W/W_i \cong W\cdot\varpi_i^\vee$. □

2.3 Lakshmibai-Seshadri Paths

We examine in detail the path crystals of $V(\varpi_i)$ where $\varpi_i^\vee$ is minuscule. For any dominant weight λ, the most canonical choice of dominant path is the straight-line path from 0 to λ, denoted $\pi = (\lambda)$. The corresponding path crystal $\mathcal{B}(\lambda)$ can be described non-recursively by the *Lakshmibai-Seshadri (LS) chains* [15].

These are saturated chains in the parabolic Bruhat order on $W\cdot\lambda$, weighted with certain rational numbers:

$$(\tau_1 \gtrdot \cdots \gtrdot \tau_m;\ 0{<}a_1{\leq}\cdots{\leq}a_{m-1}{<}1)$$

with $\tau_j{\in}W\cdot\lambda$, $a_j{\in}\mathbb{Q}$, and $m{\geq}1$. We require that if $\tau_{j+1} = \tau_j - d_j\alpha$ for $\alpha{\in}\Delta_+$, $d_j{\in}\mathbb{N}$, then $a_j = n_j/d_j$ for some $n_j{\in}\mathbb{N}$. An LS chain corresponds to the *LS path* defined as:

$$\pi = (a_1\tau_1 \star (a_2{-}a_1)\tau_2 \star (a_3{-}a_2)\tau_3 \star \cdots \star (1{-}a_{m-1})\tau_m)\,.$$

Notice that if $a_{j+1} = a_j$ then we may omit the step $0\tau_j$; in this case there may be more than one LS chain producing the same LS path. However, $B(\lambda)$ is equal to the set of all distinct LS paths [15].

Proposition 7 *Let $\varpi_i^\vee, \varpi_l^\vee$ be two minuscule coweights (possibly identical), and let $\bar{\sigma}_l$ be the linear mapping of $\mathfrak{h}_\mathbb{R}^*$ corresponding to $\varpi_l^\vee$. For each $\pi \in \mathcal{B}(\varpi_i)$, we have $\bar{\sigma}_l\pi \in \mathcal{B}(\varpi_i)$. That is, the linear mapping $\bar{\sigma}_l$ permutes the paths in $\mathcal{B}(\varpi_i)$.*

Proof. Consider an LS chain $(\tau_1{\gtrdot}\cdots{\gtrdot}\tau_m;\ 0{<}a_1{\leq}\cdots{\leq}a_{m-1}{<}1)$ corresponding to a path $\pi \in \mathcal{B}(\varpi_i)$. If $\mathfrak{g}$ is simply laced, then ϖ_i is minuscule by Lemma 5(i), and the denominator of a_j is:

$$d_j = \langle \tau_j, \alpha^\vee\rangle = \langle w\varpi_i, \alpha^\vee\rangle = \langle \varpi_i, w\alpha^\vee\rangle = 0 \text{ or } 1\,.$$

Since $0{<}a_j{=}\frac{n_j}{d_j}{<}1$, this means that $m = 1$ and $\pi = (\tau_1) = (w\varpi_i)$ for $w \in W$, a straight-line path of extremal weight. Since $\bar{\sigma}_l$ is an automorphism of the root system Δ, it permutes the elements of a W-orbit, and hence $\bar{\sigma}_l\pi$ is another straight-line LS path.

If $\mathfrak{g}$ is not simply laced, the paths of $\mathcal{B}(\varpi_i)$ are more complicated. By Lemma 6, we may assume that $\tau_{j+1} = \tau_j - d_j\alpha_{k(j)}$, where $k(j) \in \{1, \ldots, r\}$ and $\alpha_{k(j)}$ is a *simple* root. The turned path is:

$$\bar{\sigma}_l\pi = (a_1\bar{\sigma}_l\tau_1 \star (a_2{-}a_1)\bar{\sigma}_l\tau_2 \star \cdots)\,.$$

Suppose $k(j) = l$ for some j. By Lemma 5(ii), $\alpha_l^\vee$ is a short root of $\Delta^\vee$, and so is $w\alpha_l^\vee$, so we have:

$$d_j = \langle \tau_j, \alpha_l^\vee\rangle = \langle w\varpi_i^\vee, \alpha_l^\vee\rangle = \langle \varpi_i, w\alpha_l^\vee\rangle = 0 \text{ or } 1$$

by the same Lemma. This again means that $m = 1$ and $\bar{\sigma}_l\pi$ is a straight-line LS path.

Finally, suppose $k(j){\neq}l$ for all $j = 1, \ldots, m$. Then:

$$\bar{\sigma}_l\tau_{j+1} = \bar{\sigma}_l\tau_j - d_j\bar{\sigma}_l\alpha_{k(j)} = \bar{\sigma}_l\tau_j - d_j\alpha_p\,,$$

where $p := \sigma_l(k(j)) \in \{1, \ldots, r\}$. Hence $\bar{\sigma}_l\tau_j \gtrdot s_p\bar{\sigma}_l\tau_j = \bar{\sigma}_l\tau_{j+1}$, and $\bar{\sigma}_l\pi$ is an LS path. □

Although we do not need it here, we note that for any minuscule $\varpi_i^\vee$, the LS-paths of $B(\varpi_i)$ have at most two linear pieces (cf. [12]).

2.4 Twisted Demazure operators

For a Weyl group element with reduced decomposition $z = s_{i_1}\cdots s_{i_m} \in \widehat{W}$ and any path π (not necessarily dominant), we define the *crystal Demazure operator*:

$$\hat{\mathcal{B}}_z(\pi) := \{f_{i_1}^{k_1}\cdots f_{i_m}^{k_m}\pi_\lambda \mid k_1,\ldots,k_m \geq 0\}\,.$$

We can extend $\hat{\mathcal{B}}_z$ to an operator taking any set of paths Π to a larger set of paths: $\hat{\mathcal{B}}_z(\Pi) := \bigcup_{\pi\in\Pi}\hat{\mathcal{B}}_z(\pi)$. We have $\hat{\mathcal{B}}_y(\hat{\mathcal{B}}_z(\Pi)) = \hat{\mathcal{B}}_{yz}(\Pi)$ whenever $l(yz) = l(y)+l(z)$. Similarly, we let $\mathcal{B}(\Pi)$ be the set of all paths generated from Π by $f_1,\ldots,f_r$, $e_1,\ldots,e_r$.

It will be convenient to define a Demazure module $\hat{V}_z(\Lambda)$ for any $z \in \widetilde{W} = \Sigma \ltimes \widehat{W}$. Now, $\sigma \in \Sigma$ induces an automorphism of $\widehat{\mathfrak{g}}$, so for a module $\hat{V}$ we have the twisted module $\sigma\hat{V}$ defined by the action $g\odot v := \sigma^{-1}(g)v$ for $g \in \widehat{\mathfrak{g}}$, $v \in \hat{V}$. That is, $\sigma\hat{V}(\Lambda) \cong \hat{V}(\sigma\Lambda)$, and in particular $\sigma\hat{V}(\Lambda_i) \cong \hat{V}(\Lambda_{\sigma(i)})$. Now, for $z = \sigma y$ with $y \in \widehat{W}$ define: $\hat{V}_z(\Lambda) := \sigma(\hat{V}_y(\Lambda)) \subset \sigma\hat{V}(\Lambda)$, a twist of an ordinary Demazure module. Thus, $\hat{V}_{\sigma y}(\Lambda) \cong \hat{V}_{\sigma y\sigma^{-1}}(\sigma\Lambda)$ and

$$\hat{V}_{y\sigma}(\Lambda) \cong \hat{V}_y(\sigma\Lambda)\,.$$

Furthermore, $\hat{V}_{\lambda^\vee}(\Lambda_0) := \hat{V}_z(\Lambda_0)$ for $z = t_{-\nu(\lambda^\vee)}$ is a $\mathfrak{g}$-module for any dominant integral coweight $\lambda^\vee \in \oplus_{i=1}^r \mathbb{N}\varpi_i^\vee$.

The combinatorial counterpart of this construction is:

$$\hat{\mathcal{B}}_{\sigma y}(\pi) := \sigma\hat{\mathcal{B}}_y(\pi)$$

for $\sigma \in \Sigma$ and $y \in \widehat{W}$. All of our statements regarding $\hat{V}_z(\Lambda)$ and $\hat{\mathcal{B}}_z$ for $z\in\widehat{W}$ remain valid for $z\in\widetilde{W}$.

2.5 Proof of Theorems 1 and 3

Let $\lambda^\vee$ be a dominant integral coweight (not necessarily in the root lattice) which can be written:

$$\lambda^\vee = \lambda_1^\vee + \cdots + \lambda_m^\vee\,,$$

where $\lambda_j^\vee \in \{\varpi_1,\ldots,\varpi_r\}$ are minuscule fundamental coweights (not necessarily distinct) with corresponding weights λ_j and dual weights $\lambda_j^* = -w_0(\lambda_j)$. Then we claim (extending Theorem 1 to the coweight lattice) that there is an isomorphism of $\mathfrak{g}$-modules:

$$\hat{V}_{\lambda^\vee}(\Lambda_0) \cong V(\lambda_1^*)\otimes\cdots\otimes V(\lambda_m^*)\,.$$

This follows immediately from Littelmann's Character and Restriction Theorems combined with the following extension of Theorem 3.

We will choose a certain $\widehat{\mathfrak{g}}$-dominant path π_m of weight Λ_0 and show that:

$$\mathcal{B}_{\lambda^\vee}(\pi_m) = \Lambda_0 \star \mathcal{B}(\lambda_1^*)\star\cdots\star\mathcal{B}(\lambda_m^*)\,.$$

Let $\sigma_j \in \Sigma$ correspond to $\lambda_j^\vee$ for $j = 1, \dots, m$. We define π_m inductively as the last of a sequence of paths $\pi_0, \pi_1, \dots, \pi_m$:

$$\pi_0 := \Lambda_0, \qquad \pi_j := \sigma_j^{-1}(\pi_{j-1} \star \lambda_j^*)\,.$$

We may picture π_m as jumping up to level Λ_0, winding horizontally around the fundamental alcove $A = (\mathfrak{h}_\mathbb{R}^* + \Lambda_0) \cap \hat{C}$, and ending at Λ_0. Indeed, note that $\mathrm{wt}(\pi_0) = \Lambda_0$. For $j > 0$, write $\varpi := \lambda_j$ and $\Lambda := \varpi + \Lambda_0$, so that $\sigma_j(\Lambda) = \Lambda_0$ and $\sigma_j(\varpi) = -\varpi^*$. Then by induction:

$$\begin{aligned}\mathrm{wt}(\pi_j) &= \sigma_j^{-1}(\mathrm{wt}(\pi_{j-1}) + \varpi^*)\\ &= \sigma_j^{-1}\Lambda_0 + \sigma_j^{-1}\varpi^*\\ &= \Lambda - \varpi = \Lambda_0\,,\end{aligned}$$

so that each π_j has weight Λ_0. Furthermore, since $\Lambda_0 + \lambda_j^* \in \hat{C}$ and σ_j is a automorphism of $\hat{C}$, it is clear that each π_j is indeed a $\widehat{\mathfrak{g}}$-dominant path.

We will now prove Theorem 3 by showing that the Demazure operator $\mathcal{B}_{\lambda^\vee}$ "unwinds" π_m starting from its endpoint. To compute:

$$\mathcal{B}_{\lambda^\vee}(\pi_m) = \mathcal{B}_{\lambda_1^\vee} \cdots \mathcal{B}_{\lambda_m^\vee}(\pi_m)\,,$$

it suffices to prove:

Lemma 8 *For $j = m, m{-}1, \dots, 1$, we have:*

$$\begin{aligned}&\mathcal{B}_{\lambda_j^\vee}\left(\pi_j \star \mathcal{B}(\lambda_{j+1}^*) \star \cdots \star \mathcal{B}(\lambda_m^*)\right)\\ &\qquad = \pi_{j-1} \star \mathcal{B}(\lambda_j^*) \star \mathcal{B}(\lambda_{j+1}^*) \star \cdots \star \mathcal{B}(\lambda_m^*)\,.\end{aligned}$$

Proof. For $j = m$, we compute:

$$\begin{aligned}\mathcal{B}_{\lambda_m^\vee}(\pi_m) &= \hat{\mathcal{B}}_{w_m w_0 \sigma_m}\left(\sigma_m^{-1}(\pi_{m-1} \star \lambda_m^*)\right)\\ &= \hat{\mathcal{B}}_{w_m w_0}\left(\pi_{m-1} \star \lambda_m^*\right)\\ &\overset{\text{(I)}}{=} \hat{\mathcal{B}}_{w_0 w_m^*}\left(\pi_{m-1} \star \lambda_m^*\right)\\ &\overset{\text{(II)}}{=} \pi_{m-1} \star \hat{\mathcal{B}}_{w_m w_0}(\lambda_m^*)\\ &\overset{\text{(III)}}{=} \pi_{m-1} \star \hat{\mathcal{B}}_{w_0}(\lambda_m^*)\\ &= \pi_{m-1} \star \mathcal{B}(\lambda_m^*)\end{aligned}$$

The equalities are justified as follows: (I) Here $w_m^* := w_0 w_m w_0$, the longest element in $\mathrm{Stab}_W(\lambda_m^*) = \mathrm{Stab}_W(-w_0\,\lambda_m)$. (II) We say a path π is *i-neutral* if $\langle \pi(t), \alpha_i^\vee \rangle \geq 0$ and $\langle \mathrm{wt}\,\pi, \alpha_i^\vee \rangle = 0$. It is clear from the definition of f_i that $f_i(\pi \star \pi') = \pi \star f_i(\pi')$ for any path π', and the two sides are both defined or both undefined. Now note that the π_j are i-neutral for $i{=}1, \dots, r$. (III) Follows from $\hat{\mathcal{B}}_{w_0 w_m^*}(\lambda^*) = \hat{\mathcal{B}}_{w_0 w_m^*} \hat{\mathcal{B}}_{w_m^*}(\lambda^*) = \hat{\mathcal{B}}_{w_0}(\lambda^*)$.

For $j < m$, letting $\mathcal{B}_{j+1} := \mathcal{B}(\lambda_{j+1}^*) \star \cdots \star \mathcal{B}(\lambda_m^*)$, we have:

$$\begin{aligned}
\mathcal{B}_{\lambda_j^\vee}(\pi_j \star \mathcal{B}_{j+1}) &= \hat{\mathcal{B}}_{w_0 w_j^* \sigma_j}\left(\sigma_j^{-1}(\pi_{j-1} \star \lambda_j^*) \star \mathcal{B}_{j+1}\right) \\
&\overset{\text{(I)}}{=} \hat{\mathcal{B}}_{w_0 w_j^*}\left(\pi_{j-1} \star \lambda_j^* \star \sigma_j \mathcal{B}_{j+1}\right) \\
&\overset{\text{(IV)}}{=} \hat{\mathcal{B}}_{w_0 w_j^*}\left(\pi_{j-1} \star \lambda_j^* \star \mathcal{B}_{j+1}\right) \\
&\overset{\text{(II)}}{=} \pi_{j-1} \star \hat{\mathcal{B}}_{w_0 w_j^*}\left(\lambda_j^* \star \mathcal{B}_{j+1}\right) \\
&\overset{\text{(V)}}{=} \pi_{j-1} \star \mathcal{B}\left(\lambda_j^* \star \mathcal{B}_{j+1}\right) \\
&\overset{\text{(VI)}}{=} \pi_{j-1} \star \mathcal{B}(\lambda_j^*) \star \mathcal{B}_{j+1}
\end{aligned}$$

Here (I) and (II) are as above. (IV) Follows from Proposition 7. (V) Follows from [13, Prop. 12], which implies that $\lambda_j^* \star \mathcal{B}_{j+1}$ is isomorphic to a union of Demazure crystals $\mathcal{B}_y(\mu)$ with $y \geq w_m^*$. (VI) Both sides are stable under the f_i, e_i for $i = 1, \ldots, r$, and they contain the same $\mathfrak{g}$-dominant paths, hence they are identical.

This concludes the proof of the Lemma, and hence of Theorem 3. $\square$

3 Semi-infinite Crystals

3.1 Proof of Theorem 2

Fix a minuscule coweight $\varpi^\vee$ with corresponding weight ϖ, dual weight ϖ^*, and automorphism $\sigma \in \Sigma$ of order N. The N-fold concatenation $\mathcal{B}(\varpi^*) \star \cdots \star \mathcal{B}(\varpi^*)$ contains the path $\pi_N := (\sigma^{-N}(\varpi^*) \star \cdots \star \sigma^{-2}(\varpi^*) \star \sigma^{-1}(\varpi^*))$, which is $\mathfrak{g}$-dominant with weight 0. Thus $V(\varpi^*)^{\otimes N}$ possesses a corresponding invariant vector v_N.

Since $t_{-N\varpi^\vee} \in \widehat{W}$, the twisted Demazure module $V_{mN\varpi^\vee}(\Lambda_0)$ is an ordinary Demazure submodule of $\hat{V}(\Lambda_0)$. In fact, the basic $\widehat{\mathfrak{g}}$-module, considered as a $\mathfrak{g}$-module, is a direct limit of these Demazure modules:

$$\hat{V}(\Lambda_0) = \varinjlim \; V_{N\varpi^\vee} \hookrightarrow V_{2N\varpi^\vee} \hookrightarrow V_{3N\varpi^\vee} \hookrightarrow \cdots .$$

By Theorem 1, the $\mathfrak{g}$-module on the right hand side is isomorphic to:

$$\varinjlim \; V(\varpi^*)^{\otimes N} \overset{\phi_1}{\hookrightarrow} V(\varpi^*)^{\otimes 2N} \overset{\phi_2}{\hookrightarrow} V(\varpi^*)^{\otimes 3N} \overset{\phi_3}{\hookrightarrow} \cdots$$

for some injective $\mathfrak{g}$-module morphisms $\phi_1, \phi_2, \phi_3, \ldots$.

Now, if $\phi, \psi : V_1 \hookrightarrow V_2$ are any two injective $\mathfrak{g}$-module morphisms, complete reducibility implies that there exist automorphisms ξ_1, ξ_2 forming the commutative diagram:

$$\begin{tikzcd}
V_1 \arrow[r, hook, "\phi"] \arrow[d, "\xi_1"', "\wr"] & V_2 \arrow[d, "\xi_2"', "\wr"] \\
V_1 \arrow[r, hook, "\psi"] & V_2
\end{tikzcd}$$

Thus, the inclusions $\psi_m : V(\varpi^*)^{\otimes Nm} \hookrightarrow V(\varpi^*)^{\otimes N(m+1)},\ v \mapsto v_N \otimes v$ appearing in Theorem 2 define the same $\mathfrak{g}$-module as the above maps ϕ_m, and we conclude that $V(\Lambda_0)$ is isomorphic as a $\mathfrak{g}$-module to the infinite tensor product as claimed.

3.2 The Skein model

In this section we prove Theorem 4 by a crystal analog of the above argument. We introduce a notation for a path π which emphasizes the vector steps going toward the endpoint $\Lambda = \mathrm{wt}(\pi)$ rather than away from the starting point 0. Define

$$\pi = (\star v_k \star \cdots \star v_1 \vdash \Lambda) := (v' \star v_k \star \cdots \star v_1)\,,$$

the path with endpoint Λ, last step v_1, etc, and first step $v' := \Lambda - (v_k + \cdots + v_1)$, a makeweight to assure that the steps add up to Λ.

A *skein* is an infinite list:

$$\pi = (\cdots \star v_2 \star v_1 \vdash \Lambda)\,,$$

where $\Lambda \in \oplus_{i=0}^{r} \mathbb{Z}\Lambda_i$ and $v_j \in \mathfrak{h}^*_{\mathbb{R}}$ (not $\widehat{\mathfrak{h}}^*_{\mathbb{R}}$), subject to conditions (i) and (ii) below. For $i = 0, \ldots, r$ and $k > 0$, define:

$$h_i[k] := \langle\, \Lambda - (v_1 + \cdots + v_k),\ \alpha_i^\vee \,\rangle\,.$$

We require:
(i) For each i and all $k \gg 0$, we have $h_i[k] \geq 0$.
(ii) For each i, there are infinitely many k such that $h_i[k] = 0$.

We think of the skein π as a "projective limit" of the paths

$$\pi[k] := (\star v_k \star \cdots \star v_1 \vdash \Lambda) \quad \text{as} \quad k \to \infty\,.$$

The conditions on π assure that only a finite number of steps of π lie outside the fundamental chamber $\hat{C}$, and that π touches each wall of $\hat{C}$ infinitely many times. Note that π stays always at the level $\ell = \langle \Lambda, K \rangle$.

Lemma 9 *For a skein π and $i = 0, \ldots, r$, one of the following is true:*
(i) $f_i(\pi[k])$ is undefined for all $k \gg 0$;
(ii) there is a unique skein π' such that $\pi'[k] = f_i(\pi[k])$ for all $k \gg 0$.
In the second case, we define $f_i \pi := \pi'$.

Proof. Recall that a path π is i-neutral if $h_i^\pi(t) \geq 0$ for all t and $h_i^\pi(1) = 0$. For a fixed i, divide π into a concatenation: $\pi = (\cdots \star \pi_2 \star \pi_1 \star \pi_0 \vdash \Lambda)$, where each π_j is an i-neutral finite path except for π_0, which is an arbitrary finite path. Now it is clear that if $f_i(\pi_0)$ is undefined, then (i) holds. Otherwise (ii) holds and

$$f_i \pi = (\cdots \star \pi_2 \star \pi_1 \star f_i(\pi_0) \vdash \Lambda - \alpha_i\,)\,. \qquad \square$$

We can immediately carry over the definitions of the path model to skeins, including that of (Demazure) path crystals. For example, we say that π is an integral dominant skein if $\pi[k]$ is integral dominant for $k \gg 0$, and hence for all k. There exist integral dominant skeins of level $\ell = 1$ only when $\mathfrak{g}$ has a minuscule coweight. We cannot concatenate two skeins, but we can concatenate a skein π_1 and a path π_0: that is, $\pi_1 \star \pi_0 := (\,\pi_1 \star \pi_0 \vdash \mathrm{wt}(\pi_1) + \mathrm{wt}(\pi_0)\,)$.

Proposition 10 *For an integral dominant skein π of weight Λ, the crystal $\hat{\mathcal{B}}(\pi)$ is a model for $\hat{V}(\Lambda)$, and for Demazure modules.*

Proof. Given an integral dominant skein π and a Weyl group element $z \in \widetilde{W}$, we can divide $\pi = \pi_1 \star \pi_0$ in such a way that the Demazure operator $\hat{\mathcal{B}}_z$ acts on π by reflecting intervals in π_0 rather than π_1. This gives an isomorphism between the Demazure crystals generated by the path $\mathrm{wt}(\pi_1) \star \pi_0$ and by the skein π:

$$\begin{tikzcd}
\hat{\mathcal{B}}_z(\mathrm{wt}(\pi_1) \star \pi_0) \arrow[r, "\sim"] & \hat{\mathcal{B}}_z(\pi_1 \star \pi_0) = \hat{\mathcal{B}}_z(\pi) \\
\mathrm{wt}(\pi_1) \star \pi' \arrow[r, mapsto] & \pi_1 \star \pi' \quad .
\end{tikzcd}$$

This proves the assertion about Demazure modules.

Now, given an infinite chain of Weyl group elements $z_1 < z_2 < \cdots$, we have the morphisms of $\widehat{\mathfrak{g}}$-crystals:

$$\begin{tikzcd}
\hat{\mathcal{B}}_{z_1}(\Lambda) & \hat{\mathcal{B}}_{z_1}(\mathrm{wt}(\pi_1) \star \pi_0) \arrow[l, "\sim"'] \arrow[r, "\sim"] & \hat{\mathcal{B}}_{z_1}(\pi) \\
\cap & & \cap \\
\hat{\mathcal{B}}_{z_1}(\Lambda) & \hat{\mathcal{B}}_{z_2}(\mathrm{wt}(\pi_1') \star \pi_0') \arrow[l, "\sim"'] \arrow[r, "\sim"] & \hat{\mathcal{B}}_{z_2}(\pi) \\
\cap & & \cap \\
\vdots & & \vdots \\
\hat{\mathcal{B}}(\Lambda) & & \hat{\mathcal{B}}(\pi)
\end{tikzcd}$$

Since the $\widehat{\mathfrak{g}}$ crystals at the bottom are the unions of their Demazure crystals, they are isomorphic: $\hat{\mathcal{B}}(\Lambda) \cong \hat{\mathcal{B}}(\pi)$. $\square$

Now consider the situation of Theorem 4. Define the skein $\pi_\infty := (\cdots \star \pi_N \star \pi_N \vdash \Lambda_0)$, and consider the commutative diagram:

$$\begin{tikzcd}
\hat{\mathcal{B}}_{N\varpi^\vee}(\Lambda_0 \star \pi_N) \arrow[r, "\sim"] \arrow[d] & \hat{\mathcal{B}}_{\varpi^\vee}(\pi_\infty) \\
 & \cap \\
\hat{\mathcal{B}}_{2N\varpi^\vee}(\Lambda_0 \star \pi_N \star \pi_N) \arrow[r, "\sim"] \arrow[d] & \hat{\mathcal{B}}_{2N\varpi^\vee}(\pi_\infty) \\
\vdots & \cap \\
 & \vdots \\
 & \hat{\mathcal{B}}(\pi_\infty)
\end{tikzcd}$$

where the vertical maps on the left are those of the direct limit. Theorem 4 now follows from the above Proposition.

References

[1] N. Bourbaki, Groupes et Algèbres de Lie, Chapitres 4, 5 et 6, Hermann, Paris (1968).

[2] E. Date, M. Jimbo, A. Kuniba, T. Miwa and M. Okado, *A Path space realization of the basic representation of* $A_n^{(1)}$, in Infinite-Dimensional Lie Algebras and Groups, Adv. Ser. in Math. Phys. **7**, World Sci. (1989), 108–123.

[3] D. Grabiner, P. Magyar, Random walks in Weyl chambers and the decomposition of tensor powers, *J. Algebraic Combin.* **2** (1993), 239–260.

[4] J. Hong and S-J. Kang, Introduction to Quantum Groups and Crystal Bases, Grad. Stud. in Math. **42**, Amer. Math. Soc. (2002).

[5] A. Joseph, Quantum Groups and Their Primitive Ideals, Springer Verlag, New York (1994).

[6] V.G. Kac, Infinite Dimensional Lie Algebras, 3rd ed., Cambridge University Press (1990).

[7] S-J. Kang, M. Kashiwara, K.C. Misra, T. Miwa, T. Nakashima and A. Nakayashiki, Perfect crystals of quantum affine Lie algebras, *Duke Math. J.* **68** (1992), 499–607.

[8] M. Kashiwara, On crystal bases of q-analog of universal enveloping algebras, *Duke Math. J.* **63** (1991), 465–516.

[9] M. Kashiwara, On level-zero representation of quantized affine algebras, *Duke Math. J.* **112** (2002), 117–195.

[10] A. Kuniba, K. Misra, M. Okado, T. Takagi, J. Uchiyama, Demazure modules and perfect crystals, *Comm. Math. Phys.* **192** (1998), 555–567.

[11] A. Kuniba, K. Misra, M. Okado, T. Takagi, J. Uchiyama, Paths, Demazure crystals, and symmetric functions, *J. Math. Phys.* **41** (2000), 6477–6486.

[12] V. Lakshmibai and C.S. Seshadri, Standard monomial theory, Proceedings of the Hyderabad Conference on Algebraic Groups, ed. S. Ramanan, Manoj Prakashan Madras (1991), 279-323.

[13] V. Lakshmibai, P. Littelmann and P. Magyar, Standard monomial theory for Bott-Samelson varieties, *Compositio Math.* **130** (2002), 293–318.

[14] V. Lakshmibai, P. Littelmann, P. Magyar, Standard monomial theory and applications, in Representation Theories and Geometry, ed. A. Broer, Kluwer Academic Publishers (1998), 319–364.

[15] P. Littelmann, A Littlewood-Richardson formula for symmetrizable Kac-Moody algebras, *Invent. Math.* **116** (1994), 329–346.

[16] P. Littelmann, Paths and root operators in representation theory, *Ann. Math.* **142** (1995), 499–525.

[17] G. Lusztig, Introduction to Quantum groups, Progress in Mathematics **110**, Birkhäuser Verlag (1993).

[18] I.G. Macdonald, *Affine Hecke algebras and orthogonal polynomials*, Cambridge Tracts in Mathematics **157**, Cambridge University Press (2003).

[19] M. Okado, A. Schilling, M. Shimozono, Crystal bases and q-identities, in *q-series with applications to combinatorics, number theory, and physics*, Contemp. Math. **291**, Amer. Math. Soc. (2001), 29–53.

[20] G. Pappas and M. Rapoport, Local models in the ramified case I: the EL-case, *J.Alg.Geom* **12** (2003), 107–145.

[21] A. Pressley and G. Segal, Loop Groups, Oxford Univ. Press (1986).

[22] J. Stembridge, Minuscule elements of Weyl groups, *J. Algebra* **235** (2001), 722–743